Proof of a recent conjecture of Z.-W. Sun

Song Guo and Victor J. W. Guo[1]

School of Mathematical Sciences, Huaiyin Normal University, Huai'an, Jiangsu 223300, People's Republic of China
guosong77@hytc.edu.cn, jwguo@hytc.edu.cn

arXiv:1604.05019v1 [math.NT] 18 Apr 2016

Abstract. The polynomials $d_n(x)$ are defined by

$$d_n(x)=\sum_{k=0}^{n}\binom{n}{k}\binom{x}{k}2^k.$$

We prove that, for any prime p, the following congruences hold modulo p:

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(-\frac{1}{4}\right)^2\equiv\begin{cases}2(-1)^{\frac{p-1}{4}}x, & \text{if } p=x^2+y^2 \text{ with } x\equiv 1 \pmod 4,\\ 0, & \text{if } p\equiv 3 \pmod 4,\end{cases}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(-\frac{1}{6}\right)^2\equiv 0, \quad \text{if } p>3,$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(\frac{1}{4}\right)^2\equiv\begin{cases}0, & \text{if } p\equiv 1 \pmod 4,\\ (-1)^{\frac{p+1}{4}}\binom{\frac{p-1}{2}}{\frac{p-3}{4}}, & \text{if } p\equiv 3 \pmod 4.\end{cases}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(\frac{1}{6}\right)^2\equiv 0, \quad \text{if } p>5.$$

The $p\equiv 3 \pmod 4$ case of the first one confirms a conjecture of Z.-W. Sun, while the second one confirms a special case of another conjecture of Z.-W. Sun.

Keywords: Delannoy number; congurence; Fermat's little theorem

MR Subject Classifications: 11A07, 11B65, 05A10

1 Introduction

It is well known that the *Delannoy number*

$$\sum_{k=0}^{n}\binom{n}{k}\binom{m}{k}2^k=\sum_{k=0}^{n}\binom{n}{k}\binom{n+m-k}{n}$$

[1]Corresponding author.

counts lattice paths from $(0,0)$ to (m,n) using only single steps east $(1,0)$, north $(0,1)$, or northeast $(1,1)$. Recently, Z.-W. Sun [5] introduced the following polynomials

$$d_n(x)=\sum_{k=0}^{n}\binom{n}{k}\binom{x}{k}2^k.$$

and established some interesting supercongruences involving $d_n(x)$, such as

$$\sum_{k=0}^{p-1}(-1)^k d_k(x)^2\equiv(-1)^{\langle x\rangle_p}\pmod{p^2},$$

where p is an odd prime and $\langle x\rangle_p$ denotes the least non-negative integer r with $r\equiv x\pmod{p}$. He also made several interesting conjectures on congruences involving $d_n(x)$, such as (see [5, Conecture 6.2])

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(-\frac{1}{6}\right)^2\equiv\frac{p}{3}\left(\frac{p}{3}\right)\left(4\left(\frac{-2}{p}\right)-1\right)\pmod{p^2},\tag{1.1}$$

where $p>3$ is a prime and $(\frac{\cdot}{p})$ is the Legendre symbol.

In this paper, we shall prove the following result.

Theorem 1.1 *Let p be an odd prime. Then modulo p,*

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(-\frac{1}{4}\right)^2\equiv\begin{cases}2(-1)^{\frac{p-1}{4}}x, & \text{if } p=x^2+y^2 \text{ with } x\equiv 1\pmod 4,\\ 0, & \text{if } p\equiv 3\pmod 4,\end{cases}\tag{1.2}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(-\frac{1}{6}\right)^2\equiv 0,\quad \text{if } p>3,\tag{1.3}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(\frac{1}{4}\right)^2\equiv\begin{cases}0, & \text{if } p\equiv 1\pmod 4,\\ (-1)^{\frac{p+1}{4}}\binom{\frac{p-1}{2}}{\frac{p-3}{4}}, & \text{if } p\equiv 3\pmod 4.\end{cases}\tag{1.4}$$

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{4^k}d_k\left(\frac{1}{6}\right)^2\equiv 0,\quad \text{if } p>5.\tag{1.5}$$

The $p\equiv 3$ case of (1.2) was originally conjectured by Z.-W. Sun (see [5, Conjecture 6.3]), and the congruence (1.3) confirms the congruence (1.1) modulo p.

2 Proof of Theorem 1.1

In a previous paper, Guo [2, Lemma 3.1] gives the following identity:

$$d_n(x)^2=\sum_{k=0}^{n}\binom{n+k}{2k}\binom{x}{k}\binom{x+k}{k}4^k,$$

which is a special case of [4, p. 80, (2.5.32)] by noticing [4, p. 31, (1.7.1.3)] (pointed out by Wadim Zudilin).

For any odd prime p and $0 \leqslant k \leqslant p-1$, it is easy to see that

$$\frac{\binom{2k}{k}}{4^k} \equiv \begin{cases} (-1)^k \binom{\frac{p-1}{2}}{k} \pmod{p}, & \text{if } 0 \leqslant k \leqslant \frac{p-1}{2}, \\ 0 \pmod{p}, & \text{if } \frac{p+1}{2} \leqslant k \leqslant p-1. \end{cases}$$

It follows that, for any p-adic integer x,

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}}{4^k} d_k(x)^2 &\equiv \sum_{k=0}^{\frac{p-1}{2}} (-1)^k \binom{\frac{p-1}{2}}{k} \sum_{j=0}^{k} \binom{k+j}{2j} \binom{x}{j} \binom{x+j}{j} 4^j \\ &= (-1)^{\frac{p-1}{2}} \sum_{j=0}^{\frac{p-1}{2}} \binom{x}{j} \binom{x+j}{j} \binom{j}{\frac{p-1}{2}-j} 4^j \pmod{p} \end{aligned} \tag{2.1}$$

by noticing the Chu-Vandermonde identity

$$\sum_{k=j}^{\frac{p-1}{2}} (-1)^k \binom{\frac{p-1}{2}}{k} \binom{k+j}{2j} = (-1)^{\frac{p-1}{2}} \binom{j}{\frac{p-1}{2}-j}.$$

It is not difficult to see that

$$\binom{-\frac{1}{4}}{j} \binom{-\frac{1}{4}+j}{j} = (-1)^j \frac{\binom{4j}{2j}\binom{2j}{j}}{64^j} \equiv 0 \quad \text{for } \frac{p}{4} \leqslant j \leqslant \frac{p-1}{2},$$

$$\binom{-\frac{1}{6}}{j} \binom{-\frac{1}{6}+j}{j} = (-1)^j \frac{\binom{6j}{3j}\binom{3j}{j}}{432^j} \equiv 0 \quad \text{for } \frac{p}{6} \leqslant j \leqslant \frac{p-1}{2},$$

$$\binom{\frac{1}{4}}{j} \binom{\frac{1}{4}+j}{j} = (-1)^{j-1} \frac{(4j+1)\binom{4j}{2j}\binom{2j}{j}}{(4j-1)64^j} \equiv 0 \quad \text{for } j = \frac{p-1}{4} \text{ or } \frac{p+3}{4} \leqslant j \leqslant \frac{p-1}{2},$$

$$\binom{\frac{1}{6}}{j} \binom{\frac{1}{6}+j}{j} = (-1)^{j-1} \frac{(6j+1)\binom{6j}{3j}\binom{3j}{j}}{(6j-1)432^j} \equiv 0 \quad \text{for } \frac{p+3}{6} \leqslant j \leqslant \frac{p-1}{2},$$

and $\binom{j}{\frac{p-1}{2}-j} = 0$ for $0 \leqslant j < \frac{p-1}{4}$. Letting $x = \mp\frac{1}{6}$ in (2.1), we immediately obtain (1.3) and (1.5). Letting $x = \mp\frac{1}{4}$ in (2.1), we get

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}}{4^k} d_k \left(-\frac{1}{4}\right)^2 \equiv \begin{cases} (-1)^{\frac{p-1}{4}} \frac{\binom{p-1}{\frac{p-1}{2}}\binom{\frac{p-1}{2}}{\frac{p-1}{4}}}{16^{\frac{p-1}{4}}} \pmod{p}, & \text{if } p \equiv 1 \pmod 4, \\ 0 \pmod{p}, & \text{if } p \equiv 3 \pmod 4, \end{cases}$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}}{4^k} d_k \left(\frac{1}{4}\right)^2 \equiv \begin{cases} 0 \pmod{p}, & \text{if } p \equiv 1 \pmod 4, \\ (-1)^{\frac{p+1}{4}} \frac{(4p+2)\binom{p+1}{\frac{p+1}{2}}\binom{\frac{p+1}{2}}{\frac{p+1}{4}}}{4p \cdot 16^{\frac{p+1}{4}}} \pmod{p}, & \text{if } p \equiv 3 \pmod 4, \end{cases}$$

We now suppose that p is a prime such that $p\equiv 1 \pmod 4$ and $p=x^2+y^2$ with $x\equiv 1 \pmod 4$. Then by the Beukers-Chowla-Dwork-Evans congruence [1,3]:

$$\binom{\frac{p-1}{2}}{\frac{p-1}{4}} \equiv \frac{2^{p-1}+1}{2}\left(2x-\frac{p}{2x}\right) \pmod{p^2},$$

and Fermat's little theorem, we have $\binom{\frac{p-1}{2}}{\frac{p-1}{4}} \equiv 2x \pmod p$. Moreover, we have

$$\binom{p-1}{\frac{p-1}{2}} \equiv 16^{\frac{p-1}{4}} \equiv 1 \pmod p.$$

This proves (1.2). Finally, suppose that p is a prime with $p\equiv 3 \pmod 4$. Then

$$\frac{(4p+2)\binom{p+1}{\frac{p+1}{2}}\binom{\frac{p+1}{2}}{\frac{p+1}{4}}}{4p\cdot 16^{\frac{p+1}{4}}} \equiv \binom{\frac{p-1}{2}}{\frac{p-3}{4}} \pmod p.$$

This proves (1.4).

Acknowledgment. This work was partially supported by the National Natural Science Foundation of China (grant no. 11371144).

References

[1] S. Chowla, B. Dwork, and R. Evans, On the mod p^2 determination of $\binom{(p-1)/2}{(p-1)/4}$, J. Number Theory 24 (1986), 188–196.

[2] V.J.W. Guo, Proof of Sun's conjectures on integer-valued polynomials, preprint 2016, arXiv:1601.04250.

[3] H. Pan, An elementary approach to $\binom{(p-1)/2}{(p-1)/4}$ modulo p^2, Taiwanese J. Math. 16 (2012), 2197–2202.

[4] L.J. Slater, Generalized Hypergeometric Functions, Cambridge University Press, Cambridge, 1966.

[5] Z.-W. Sun, Supercongruences involving dual sequences, preprint, 2015, arXiv:1512.00712v3.